

ASYMPTOTIC PROPERTIES OF STOCHASTIC POPULATION DYNAMICS

Sulin Pang¹, Feiqi Deng² and Xuerong Mao³

¹Department of Accountancy & Institute of Finance Engineering,
School of Management, Jinan University, Guangzhou 510632, P.R. China

²Systems Engineering Institute,
South China University of Technology, Guangzhou 510640, China

³ Department of Statistics and Modelling Science,
University of Strathclyde, Glasgow G1 1XH, U.K.

Abstract. In this paper we stochastically perturb the classical Lotka–Volterra model

$$\dot{x}(t) = \text{diag}(x_1(t), \dots, x_n(t))[b + Ax(t)]$$

into the stochastic differential equation

$$dx(t) = \text{diag}(x_1(t), \dots, x_n(t))[(b + Ax(t))dt + \beta dw(t)].$$

The main aim is to study the asymptotic properties of the solution. It is known (see e.g. [3, 20]) if the noise is too large then the population may become extinct with probability one. Our main aim here is to find out what happens if the noise is relatively small. In this paper we will establish some new asymptotic properties for the moments as well as for the sample paths of the solution. In particular, we will discuss the limit of the average in time of the sample paths.

Keywords. Brownian motion, stochastic differential equation, Itô’s formula, average in time, boundedness.

AMS (MOS) subject classification: 60J65, 60H10, 34K40.

1 Introduction

The classical Lotka–Volterra model for n interacting species is described by the n -dimensional differential equation

$$\frac{dx(t)}{dt} = \text{diag}(x_1(t), \dots, x_n(t))[b + Ax(t)], \quad (1.1)$$

where

$$x = (x_1, \dots, x_n)^T, \quad b = (b_1, \dots, b_n)^T, \quad A = (a_{ij})_{n \times n}.$$

There is an extensive literature concerned with the dynamics of this model and we here only mention Ahmad and Rao [1], Bereketoglu and Gyori [4], Freedman and Ruan [9], He and Gopalsamy [11], Kuang and Smith [15], Teng